

# MARINE SCIENCE AND TECHNOLOGY BULLETIN

## RESEARCH ARTICLE

### Effect of *Tilia tomentosa* methanolic extract on growth performance, digestive enzyme activity, immune response and haematological indices of common carp (*Cyprinus carpio*).

Ahmed Alhadi Almabrok<sup>1,2</sup>, Iman Daw Amhamed<sup>1,2</sup>, Gamaia Ali Mohamed<sup>1,2</sup>,  
Soner Bilen<sup>2</sup>, Tarek Abdalsalam Salem Altief<sup>1,2\*</sup>

<sup>1</sup> Kastamonu University, Institute of Science, Department of Aquaculture, Kastamonu, Turkey

<sup>2</sup> Kastamonu University, Faculty of Fisheries and Aquaculture, Department of Aquaculture Kastamonu, Turkey

#### ARTICLE INFO

##### Article History:

Received: 04.05.2018  
Received in revised form: 23.05.2018  
Accepted: 23.05.2018  
Available online: 30.06.2018

##### Keywords:

*Tilia tomentosa*  
Common carp  
Growth  
Haematology  
Digestive enzyme activity  
Immune indices

#### ABSTRACT

This study was conducted to determine the effect of dietary supplementation with *Tilia tomentosa* on the growth performance, digestive enzyme activity, haematological indices and nonspecific immune indices of juvenile common carp (*Cyprinus carpio*). Fish with an average weight of  $4.35 \pm 0.16$  g were fed a diet supplemented with an aqueous methanolic extract of *T. tomentosa* at a dose of 0% (control), 0.01%, 0.05% or 0.1% for 45 days. The final weight, weight gain and specific growth rate were observed to be significantly higher for the 0.01% and 0.1% groups compared with the control group ( $P < 0.05$ ). The feed conversion ratio was significantly decreased in the 0.05% and 0.1% groups compared with the control ( $P < 0.05$ ). The activities of various digestive enzymes (amylase, lipase and trypsin) were also measured and no significant differences were observed compared to the control ( $P > 0.05$ ). The mean cell volume of the 0.01% group was significantly increased compared to the control ( $P < 0.05$ ) and increased lysozyme activity was observed in the 0.05% and 0.1% groups. Respiratory burst activity was significantly increased ( $P < 0.05$ ) on days 15 and 30 for the 0.1% and 0.05% groups, respectively. No differences were observed for myeloperoxidase activity among the four groups. These results suggest that aqueous methanolic extract of *T. tomentosa* has a growth-promoting and immunostimulatory effect on common carp.

#### Please cite this paper as follows:

Almabrok, A.A., Amhamed, I.D., Mohamed, G.A., Bilen, S., Altief, T.A.S. (2018). Effect of *Tilia tomentosa* methanolic extract on growth performance, digestive enzyme activity, immune response and haematological indices of common carp (*Cyprinus carpio*). *Marine Science and Technology Bulletin*, 7(1): 12-20.

#### Introduction

Aquaculture is the most important sector in the world.

Also it considered as the most growing food produce industry with an average growth rate more than 7.7% per year through the last decades, the majority of aquaculture production are come from Asia (Gjedrem et al., 2012). Approximately 600 aquatic species are raised in captivity in around 190 countries for produce in fish culture system of varying input intensity

\* Corresponding author  
E-mail address: telhasy@yahoo.com (T.A.S. Altief)

and technological sophistication (Aklakur et al., 2015).

During last decades there has been a continuous growth of aquaculture industries all over the world and such intensive production would experience disease problems. Infectious diseases that occur as sporadic events in wild fish populations may cause high mortalities when appearing in intensive fish farming (Gudding et al., 1999). Enhancing the immune system of cultured fish appears to be the most promising method for preventing disease. Fish typically depend on nonspecific immune mechanisms to a much greater extent than most animals (Chakrabarti and Vasudeva, 2006).

Medicinal plants contain phytochemicals, which exhibit biological activities such as the prevention of chronic degenerative diseases (Fukumoto and Mazza, 2000). Herbal medicines have been used in aquaculture to promote growth, improve the immune system, and stimulate the appetite and fight against microbes and other stressors, due to the presence of various active compounds like flavanoids, alkaloids, phenolics, pigments, steroids, terpenoids and essential oils (Citarasu, 2010). Recently, immunostimulants of herbal origin have been shown to boost the disease resistance of fish towards a number of diseases by improving their nonspecific and specific defence mechanisms (Harikrishnan et al. 2011). Indeed, many herbs have been reported to enhance the immune response of fish (Khondoker et al, 2016).

The *Tilia tomentosa* extracts contained flavonols (quercetin, kaempferol, apigenin derivatives) as principal components with the exception of a single commercial extract with hydroxycinnamic acids as the most abundant metabolites (İleri et al., 2015). *T. tomentosa* is used as a medicinal plant in Turkish folk medicine (Baser et al., 2005). The present study was carried out to determine whether the *T. tomentosa* extract influences the digestive enzyme activity, growth, haematological parameters, or activity of the nonspecific immune response of the common carp (*Cyprinus carpio*).

## Material and Methods

### Preparation of *T. tomentosa* Extract

The plants were collected from the Kastamonu province in the north of Turkey and extracted using a methanol extraction method (Pakravan et al., 2012) with some modifications (Bilen et al., 2016) as follows: All ripe parts of the fruit except the seeds were ground in a mechanical grinder to a fine powder. Every 50 g of the ground plant were mixed with 1 L of 40% methanol (Sigma-Aldrich) and the mixture was allowed to stand at room temperature for three days with brief shaking once a day to mix. The extract was then filtered through filter paper (Whatman filter paper No. 1) and the filtrate was collected and evaporated in a rotary evaporator at 55-65 °C to remove the alcohol from the fruit extract. The final crude product was dissolved in distilled water and kept in a flask at 4 °C for later experiments.

### Experimental Design

Common carp (*C. carpio*) with an average body weight of

4.35 ± 0.16 g were obtained from a commercial fish farm in Antalya, Turkey. The fish were transported to the Faculty of Fisheries and Aquaculture, Kastamonu University, Turkey. The 480 carp were randomly divided into four groups of three replicates each (i.e. four groups of three aquariums, 12 aquariums in total) with 40 fish per aquarium. The fish were acclimatised for two weeks before the experiment started. During acclimation, the fish were fed with a commercial diet twice a day. During the experimental period, the fish were fed the commercial diet supplemented with *T. tomentosa* extract at the following percentages (w/w): 0%, 0.01%, 0.05%, or 0.1%. The fish were fed by hand with the experimental diet for 45 days to satiation twice a day (at 9 am and 4 pm). The fish were maintained under a natural photoperiod (12-h dark/12-h light). The water quality parameters were checked daily and were within the accepted margins throughout the experiment (dissolved oxygen, 6.8-7.2 mg/L; pH, 7.7-8.5; water temperature, 25-28 °C).

### Sample Collection

Every 15 days of the feeding trial (i.e. on days 15, 30 and 45), three fish per aquarium (a total of nine carp from each experimental group) were randomly chosen, anaesthetised by phenoxyethanol at 0.01 mL/L and individually weighed and sampled. The kidney tissues were collected and transferred individually to 1.5 mL RPMI-1640 medium (Invitrogen, Carlsbad, CA, USA) for direct assay of the immunological parameters. Blood was collected on day 45 of the feeding trials from the caudal vein using heparinised syringes in EDTA-tubes, and later used for the direct assay of the haematological parameters. Samples of the intestine were also collected on day 45 of the feeding trials. These samples were cleaned to remove waste and all visible fat and then stored at -80 °C for the digestive enzyme assay.

### Growth Performance Parameters

The weight of each fish was individually measured at the beginning and end of the study. The growth performance was calculated according to the following equations (Tekinay and Davies, 2001):

$$WG(\%) = 100 \times \left[ \frac{(\text{Final Fish Weight} - \text{Initial Fish Weight})}{\text{Initial Fish Weight}} \right]$$

$$SGR(\text{weight \% d}^{-1}) = 100 \times \left[ \frac{\ln(\text{Final Fish Weight}) - \ln(\text{Initial Fish Weight})}{\text{Experimental Days}} \right]$$

$$FCR = \frac{\text{Feed Intake (g)}}{WG (g)}$$

$$SR(\%) = 100 \times \left[ \frac{\text{Final Number of Fish}}{\text{Initial Number of Fish}} \right]$$

In these formulae, WG indicates weight gain, SGR indicates specific growth rate, FCR indicates feed conversion ratio, and SR indicates survival rate.

### Digestive Enzyme Activity

The intestine samples were homogenised with a Potter Elvehjem homogeniser in cold double-distilled water (0.1 g/mL) and centrifuged at 9,000 rpm for 20 min at 4 °C. The

supernatant was removed and stored at  $-80^{\circ}\text{C}$  to test for digestive enzyme activity as follows. The amylase activity was determined using 2% starch (Sigma-Aldrich) as a substrate according to the Worthington (1991) method. The lipase activity was determined using hydrolysis of 4-nitrophenyl myristate (Sigma-Aldrich) according to the method described by Gawlicka et al. (2000). The trypsin activity was determined using the method of Erlanger et al. (1961) and benzoyl-DL-arginine p-nitroanilide (Sigma-Aldrich) as the substrate. The protein concentrations were evaluated with Bradford (1976) method.

### Haematological Profiles

White blood cell counts ( $\text{WBC} \times 10^7 \text{ mm}^{-3}$ ), red blood cell counts ( $\text{RBC} \times 10^6 \text{ mm}^{-3}$ ), haemoglobin levels (Hb, g/dL) and haematocrit measurements (Hct, %) were measured according to the methods described by Blaxhall and Daisley (1973). Blood indices included the mean cell volume (MCV, fL), mean cell Hb (MCH, pg) and mean cell Hb concentration (MCHC, %), which were calculated according to the formulae of Lewis et al. (2001).

### Nonspecific Immune Parameters

Head kidney cells were isolated from euthanised *C. carpio* according to the method of Kono et al. (2012) with slight modifications as follows. Briefly, the head kidney tissue was carefully removed and gently pushed through a 100- $\mu\text{m}$  nylon mesh (John Stanier & Co., Whitefield, Manchester, UK) into RPMI-1640 medium (Invitrogen) supplemented with 5% foetal bovine serum (Invitrogen) and a 1% solution of 10,000 g/mL streptomycin plus 10,000 U/mL penicillin (Invitrogen). This mixture was then pushed through a 40- $\mu\text{m}$  nylon mesh cell strainer (Becton, Dickinson & Co., Franklin Lakes, NJ, USA). The final homogenate was placed in a 3-mL Falcon tube. Head kidney cell suspensions were pelleted at 1,800 rpm for 3 min at  $4^{\circ}\text{C}$ . After centrifugation, the supernatant was collected to measure myeloperoxidase activity using 3,3,5,5-tetramethyl benzidine hydrochloride (Sigma-Aldrich) as the substrate (Sahoo et al., 2005). The lysozyme activity was measured using a lyophilised *Micrococcus lysodeikticus* bacterial cell solution (Sigma-Aldrich) as the substrate (Bilen et al., 2014). Each pellet was resuspended in 1 mL of the same medium to directly assay nitroblue tetrazolium (Sigma-Aldrich) reduction, according to the method described by Biswas et al. (2013).

### Statistical Analysis

The results were analysed with SPSS software. One-way ANOVA and Duncan's multiple range test were used to determine the significant differences between groups. All results were expressed as the mean  $\pm$  SE and  $P < 0.05$  was considered statistically significant.

## Results

### Growth

The final weight (FW), FCR, SGR, WG and SR of common carp fed on the experimental diets for 45 days were determined and presented in Table 1.

**Table 1.** Growth indices of *Cyprinus carpio* supplemented with a methanolic extract of *T. tomentosa* for 45 days.

| Groups  | IW (g)            | FW (g)            | WG (%)              | FCR               | SGR (%/day)       |
|---------|-------------------|-------------------|---------------------|-------------------|-------------------|
| Control | $4.09 \pm 0.10^a$ | $6.88 \pm 0.08^a$ | $168.07 \pm 5.06^a$ | $1.68 \pm 0.01^a$ | $1.15 \pm 0.04^a$ |
| 0.01%   | $4.06 \pm 0.05^a$ | $7.71 \pm 0.09^b$ | $189.87 \pm 6.28^b$ | $1.78 \pm 0.02^a$ | $1.42 \pm 0.02^b$ |
| 0.05%   | $4.23 \pm 0.06^a$ | $7.25 \pm 0.07^a$ | $171.23 \pm 4.57^c$ | $1.24 \pm 0.01^b$ | $1.20 \pm 0.02^a$ |
| 0.10%   | $4.12 \pm 0.14^a$ | $7.56 \pm 0.21^b$ | $183.22 \pm 6.19^d$ | $1.56 \pm 0.01^c$ | $1.34 \pm 0.03^b$ |

**Note:** Values represent the mean  $\pm$  SE; different superscript letters in a column indicate significant differences between groups ( $P < .05$ ). IW: initial weight, FW: final weight, WG: weight gain, FCR: feed conversion ratio, SGR: specific growth rate, SR: survival rate.

Significant increases ( $P < 0.05$ ) in FW, WG and SGR were observed for fish supplemented with 0.01% and 0.1% extract compared to the control. In addition, the FCR was significantly ( $P < 0.05$ ) lower in the 0.05% and 0.1% supplemented groups compared with the control group. No significant differences were observed between the FW, FCR, SGR and SR of the other treated groups compared with the control group.

### Digestive Enzymes

The digestive enzyme activity levels for each experimental group are shown in Table 2. No significant differences in the activity of amylase, lipase or trypsin were observed for fish supplemented with 0.01%, 0.05%, or 0.1% *T. tomentosa* extract compared with the control group.

**Table 2.** Digestive enzyme activity in the intestines of *Cyprinus carpio* supplemented with a methanolic extract of *T. tomentosa* for 45 days.

| Groups  | Enzymes         |                     |                 |
|---------|-----------------|---------------------|-----------------|
|         | Amylase         | Lipase              | Trypsin         |
| Control | $1.84 \pm 0.54$ | $0.009^a \pm 0.003$ | $0.11 \pm 0.01$ |
| 0.01%   | $1.58 \pm 0.51$ | $0.009 \pm 0.001$   | $0.08 \pm 0.01$ |
| 0.05%   | $3.95 \pm 1.26$ | $0.004 \pm 0.002$   | $0.12 \pm 0.01$ |
| 0.10%   | $3.96 \pm 1.58$ | $0.006 \pm 0.015$   | $0.08 \pm 0.05$ |

**Note:** Values represent the mean  $\pm$  SE.

### Haematological Profiles

Fish supplemented with *T. tomentosa* extract at different concentrations had the same haematological parameters as

the control group throughout the experimental periods (Table 3), with the exception of the MCV of the 0.01% group on day 30 of the experiment ( $P < 0.05$ ) and the MCHC of the control on day 30 of the experiment (highest recorded result compared with the other groups;  $P < 0.05$ ).

### Nonspecific Immune Parameters

The immunostimulatory effects of the *T. tomentosa* extract are shown in Figures 1, 2 and 3. The lysozyme activity of fish supplemented with different concentrations of *T. tomentosa* extract was increased on day 30 of the experiment; this increase was found to be significant ( $P < 0.05$ ) only for the 0.1% supplementation compared with the control (Figure 1). On the other hand, the lysozyme activity decreased significantly on day 45 of the experiment for the 0.01% supplementation compared with the control ( $P < 0.05$ ). The lysozyme activity levels of the other supplementation groups did not change significantly over time.

The myeloperoxidase activity of the treated fish did not significantly differ from that of the control fish during the


**Table 3.** Haematological profiles of *Cyprinus carpio* supplemented with a methanolic extract of *T. tomentosa* for 45 days.

| Groups               |         | WBC<br>( $\times 10^7$ cells mm <sup>-3</sup> ) | RBC<br>( $\times 10^6$ cells mm <sup>-3</sup> ) | Hb (g dl <sup>-1</sup> ) | Hct (%)      | MCV (fl)       | MCH (pg)     | MCHC (%)       |
|----------------------|---------|-------------------------------------------------|-------------------------------------------------|--------------------------|--------------|----------------|--------------|----------------|
| 15 <sup>th</sup> day | Control | 24.47 ± 0.18                                    | 1.98 ± 0.04                                     | 8.22 ± 0.07              | 27.82 ± 0.84 | 141.11 ± 7.12  | 41.57 ± 1.05 | 296.47 ± 7.63  |
|                      | 0.01%   | 25.63 ± 1.26                                    | 1.86 ± 0.09                                     | 7.77 ± 0.25              | 27.68 ± 0.55 | 148.92 ± 3.73  | 41.76 ± 0.78 | 281.13 ± 4.32  |
|                      | 0.05%   | 22.57 ± 1.48                                    | 1.97 ± 0.06                                     | 8.07 ± 0.10              | 27.88 ± 0.53 | 142.44 ± 6.20  | 41.14 ± 0.89 | 289.85 ± 8.25  |
|                      | 0.10%   | 26.03 ± 0.97                                    | 1.98 ± 0.05                                     | 8.27 ± 0.11              | 28.09 ± 0.87 | 143.29 ± 8.22  | 42.02 ± 1.08 | 295.31 ± 10.34 |
| 30 <sup>th</sup> day | Control | 22.61 ± 1.67                                    | 1.97± 0.08                                      | 8.37 ± 0.17              | 25.99 ± 0.47 | 132.72 ± 2.71  | 42.69 ± 0.89 | 321.95 ± 2.56* |
|                      | 0.01%   | 26.84 ± 0.91                                    | 1.76 ± 0.09                                     | 7.72 ± 0.15              | 26.72 ± 0.33 | 153.71 ± 6.98* | 44.23 ± 1.42 | 289.10 ± 4.66  |
|                      | 0.05%   | 26.34 ± 1.00                                    | 1.81 ± 0.03                                     | 7.68 ± 0.36              | 25.59 ± 0.73 | 141.85 ± 2.26  | 42.51 ± 1.21 | 300.12 ± 7.38  |
|                      | 0.01%   | 26.35 ± 0.45                                    | 1.93 ± 0.07                                     | 8.08 ± 0.10              | 26.84 ± 0.09 | 140.89 ± 5.95  | 42.1 2± 1.32 | 301.50 ± 4.48  |
| 45 <sup>th</sup> day | Control | 23.76 ± 1.85                                    | 1.90 ± 0.09                                     | 7.31 ± 0.33              | 25.59 ± 0.62 | 134.97 ± 4.07  | 38.45 ± 0.55 | 285.78 ± 7.19  |
|                      | 0.10%   | 24.16 ± 1.73                                    | 2.02 ± 0.05                                     | 7.81 ± 0.19              | 26.35 ± 0.64 | 131.08 ± 4.68  | 38.70 ± 1.29 | 269.27 ± 1.26  |
|                      | 0.50%   | 26.74 ± 1.13                                    | 1.97 ± 0.07                                     | 7.80 ± 0.10              | 26.18 ± 0.27 | 133.49 ± 5.52  | 39.65 ± 1.08 | 298.42 ± 6.54  |
|                      | 0.01%   | 27.41 ± 0.15                                    | 1.92 ± 0.06                                     | 7.70 ± 0.23              | 26.87 ± 0.48 | 140.42 ± 4.23  | 40.14 ± 0.30 | 286.51 ± 7.17  |

The myeloperoxidase activity of the treated fish did not significantly differ from that of the control fish during the study period (Figure 2).


The respiratory burst level of the 1% group was significantly lower ( $P < 0.05$ ) than that of the control on day 15 of the experiment (Figure 3). No significant differences in the respiratory burst level were observed among the other

study period (Figure 2).




**Figure 1.** Lysozyme activity in kidney leucocytes of *Cyprinus carpio* fed with different experimental diets for 45 days. Values are expressed as the mean  $\pm$  SE. Different symbols above the bars indicate significant differences between groups ( $P < 0.05$ ).

treated groups on day 15 compared to the control group. On day 30 of the experiment, significantly higher levels of respiratory burst were observed for the 0.05% and 0.1% groups, while a significantly lower level was observed for the 0.01% group compared with the control ( $P < 0.05$ ). The same pattern was exhibited on day 45 of the experiment ( $P < 0.05$ ).



**Figure 2.** Myeloperoxidase activity in kidney leucocytes of *Cyprinus carpio* fed with different experimental diets for 45 days. Values are expressed as the mean  $\pm$  SE.



**Figure 3.** Respiratory burst level in kidney leucocytes of *Cyprinus carpio* fed with different experimental diets for 45 days. Values are expressed as the mean  $\pm$  SE. Different letters above the bars indicate significant differences between groups ( $P < 0.05$ ).

## Discussion

In the present study, *T. tomentosa* was shown to promote the growth of common carp based on observations of increased WG, SGR and efficiency of feed conversion. These results agree with those of previous studies demonstrating that medicinal plants promote the growth of various aquatic animals (Kour et al., 2004; Kaleeswaran et al., 2011; Ojha et al., 2014). The WG was significantly improved when Japanese flounder (*Paralichthys olivaceus*) were supplemented with an herbal mixture to 500 mg/kg (Seung-Cheol et al., 2007). Most probably fat was used for energy, and protein was used for growth in the herbal-supplemented diet (Yilmaz et al., 2012). Nile tilapia fingerlings fed with a basal diet containing 0, 0.5, 1 and 1.5 g/100 g fenugreek (*Trigonella foenum-graecum*) seed meal for three months (Mostafa et al. 2009) and they found that the use of 1 g/100 g fenugreek seed meal improved the fish growth performance. However, although some herbs have positive effects on the fish growth (Xie et al., 2008; Mahdavi et al., 2013), other herbal supplements have not been observed to have any effects. For example, Farahi et al. (2012) found that dietary supplementation with

*Aloe vera* did not promote the growth performance of rainbow trout (*Oncorhynchus mykiss*). Yilmaz et al. (2012) reported that the WG, FCR and SGR of sea bass (*Dicentrarchus labrax*) were unaffected by 1000 mg/kg dietary rosemary (*Rosmarinus officinalis*) and fenugreek. Also, the growth rate of koi carp (*Cyprinus carpio*) was unaffected by dietary supplementation with tetra (*Cotinus cogglyria*) (Bilen et al., 2013).

Digestive enzymes play a significant role in the hydrolysis of proteins, lipids and carbohydrates, thereby assisting with the assimilation of nutrients. These nutrients are transported into the tissues and incorporated into cellular materials or used as an energy source for growth and reproduction (Furne et al., 2005). The digestion of foods begins with the digestive enzymes in the stomach and continues in the intestine with the digestive enzymes secreted by the pancreas, such as trypsin, chymotrypsin, amylase and lipase (Cockson and Bourne, 1972; Moriarty, 1973; Fang and Chiou, 1989).

Our study did not find any significant differences in the amylase, lipase and trypsin activity levels in any of the experimental groups compared with the control group. This result is in line with a study done by Iqbal et al. (2016), who found no significant differences in the digestive enzyme activity and haematology of juvenile *Labeo rohita* following supplementation with different plant and animal origin feeds (fishmeal). In contrast, a study by Kawai and Ikeda (1973) reported an increase in amylase activity in *O. mykiss* when fed a diet containing increased amounts of plant protein.

Many fish physiologists have concentrated on haematological studies as this has proved a valuable diagnostic approach for evaluating fish quality (Kori-Siakpre et al., 2005; Oluyemi et al., 2008; Patra et al., 2014). Variations in the haematological parameters of fish are due to environmental stress (Hickey, 1982), malnutrition (Casillas and Smith, 1977), gender (Siddique and Naseem, 1979; Collazos et al., 1998), fish size (Garcia et al., 1992), seasonal differences and breeding efficiency (Cech and Wohlschlang, 1981). The blood characteristics of fish are therefore an effective and sensitive index for monitoring physiological and pathological changes (Iwama et al., 1976; Chakrabarti and Banerjee, 1988; Orun et al., 2003; Patra et al., 2014). The present study indicated no significant effect on the blood parameters of *C. carpio* supplemented with 0.01%, 0.05%, or 0.1% *T. tomentosa* extract over the study period. This result suggested that the extract tested here did not stress the fish physiologically.

Lysozyme is an important enzyme in the humeral nonspecific defence mechanism that provides defence against microbial invasion (Evelyn, 2002). The bactericidal action of this enzyme involves the hydrolysis of the peptidoglycan layers of the bacterial cell wall, which lyses the cells and prevents colonisation by microorganisms (Saurabh and Sahoo, 2008). Lysozyme also induces antibacterial activity in the presence of a complement (Harikrishnan et al., 2011). The present study recorded significantly increased lysozyme activity in the 1% supplemented group compared with the control. Increasing lysozyme activity is in agreement with several reports on herbal immunostimulants (Rao et al., 2006;

Choi et al., 2008; Bilen et al., 2011). Similar results were observed for common carp when fed a diet supplemented with methanolic extracts of *Cotinus coggygria* (Bilen et al., 2014) or various Chinese herbal extracts (Jian and Wu, 2004).

Neutrophils contain myeloperoxidase in their cytoplasmic granules (Rodriguez et al., 2003). Myeloperoxidase is an important enzyme with microbiocidal properties as it utilises a reactive oxygen species ( $H_2O_2$ ) to produce hypochlorous acid (Dalmo et al., 1997). This process is believed to be key in killing microorganisms (Johnston, 1978). Although some studies have shown increased myeloperoxidase activity in fish following supplementation with, for example, quercetin or black cumin seed oil (Awad et al., 2013; Alexander et al., 2010; Bilen et al. 2013), our study showed no significant effect on myeloperoxidase activity following supplementation with *T. tomentosa* extract at different concentrations.

Phagocytosis and the respiratory burst response by phagocytes in blood and tissues present a major antibacterial defense mechanism in fish (Secombes, 1996). Respiratory burst activity measured by nitroblue tetrazolium (NBT) is one of the most important bactericidal mechanisms in fish (Secombes and Fletcher, 1992). In this study, the respiratory burst levels of common carp were significantly decreased on days 15, 30 and 45 of supplementation with 0.1%, 0.05% and 0.01% *T. tomentosa* extract, respectively, compared with the control group. For the groups supplemented with 0.05% and 0.1% extract, on days 30 and 45 of the experiment we found a significant increase in this parameter compared with the control. This was similar to results observed by Bilen et al. (2011) when *O. mykiss* was supplemented with *Cotinus coggygria* leaves. Harikrishnan et al. (2010) reported a significant increase in respiratory bursts for olive flounder supplemented with three different Korean plants. Haghighi and Rohani (2013) reported a significantly higher respiratory burst level in rainbow trout fed a commercial diet containing *Zingiber officinale*. Our results are also in agreement with several studies on the use of dietary immunostimulants in various fish species (Yin et al., 2009; Bilen and Bulut, 2010).

## Acknowledgement

An earlier version of this paper has been presented at International Congress on Engineering and Life Science ICCELIS 2018, Kastamonu, Turkey.

## Conflict of Interest

The authors declare that there is no conflict of interest.

## References

- Aklakur, M., Rather, M.A. & Kumar, N. (2015). Nano delivery: an emerging avenue for nutraceuticals and drug delivery. *Critical reviews in Food Science and Nutrition*, **56**(14): 2352-2361.
- Alexander, C.P., Kirubakaran, C.J.W. & Michael, R.D. (2010). Water soluble fraction of *Tinospora cordifolia* leaves enhanced the non-specific immune mechanisms and disease resistance in *Oreochromis mossambicus*. *Fish & Shellfish Immunology*, **29**(5): 765-772.
- Alishahi, M., Ranjbar, M., Ghorbanpoor, M., Peyghan, R. & Mesbah, M. (2010). Effects of dietary *Aloe vera* on some specific and nonspecific immunity in the common carp (*Cyprinus carpio*). *Iranian Journal of Veterinary Medicine*, **4**(3): 189-195.
- Awad, E., Austin, D. & Lyndon, A.R. (2013). Effect of black cumin seed oil (*Nigella sativa*) and nettle extract (Quercetin) on enhancement of immunity in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Aquaculture*, **388**: 193-197.
- Baser, K.H.C., Tümen, G., Malyer, H. & Kirimer, N. (2006). Plants used for common cold in Turkey. In *Proceedings of the IVth International Congress of Ethnobotany (ICEB 2005)*, **133**, p. 137.
- Bilen, S., Biswas, G., Otsuyama, S., Kono, T., Sakai, M. & Hikima, J.I. (2014). Inflammatory responses in the Japanese pufferfish (*Takifugu rubripes*) head kidney cells stimulated with an inflammasome-inducing agent, nigericin. *Developmental & Comparative Immunology*, **46**(2): 222-230.
- Bilen, S., Ünal, S. & Güvensoy, H. (2016). Effects of oyster mushroom (*Pleurotus ostreatus*) and nettle (*Urtica dioica*) methanolic extracts on immune responses and resistance to *Aeromonas hydrophila* in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, **454**: 90-94.
- Bilen, S. & Bulut, M. (2010). Effects of laurel (*Laurus nobilis*) on the non-specific immune responses of rainbow trout (*Oncorhynchus mykiss*, Walbaum). *Journal of Animal and Veterinary Advances*, **9**(8): 1275-1279.
- Bilen, S., Bulut, M. & Bilen, A.M. (2011). Immunostimulant effects of *Cotinus coggygria* on rainbow trout (*Oncorhynchus mykiss*). *Fish & Shellfish Immunology*, **30**(2): 451-455.
- Bilen, S., Yilmaz, S., & Bilen, A.M. (2013). Influence of tetra (*Cotinus coggygria*) extract against *Vibrio anguillarum* infection in koi carp, *Cyprinus carpio* with reference to haematological and immunological changes. *Turkish Journal of Fisheries and Aquatic Sciences*, **13**(3): 517-522.
- Bilen, S., Yilmaz, S., Bilen, A.M. & Biswas, G. (2014). Effects of Dietary Incorporation of Tetra (*Cotinus coggygria*) Extract on Immune Response and Resistance to *Aeromonas hydrophila* in Koi Carp (*Cyprinus carpio*). *The Israeli Journal of Aquaculture-Bamidgeh*, **66**: 1-6.
- Biswas, G., Korenaga, H., Nagamine, R., Kawahara, S., Takeda, S., Kikuchi, Y., Dashnyam, B., Yoshida, T. & Sakai, M. (2013). Cytokine mediated immune responses in the Japanese pufferfish (*Takifugu rubripes*) administered with heat-killed *Lactobacillus paracasei* spp. *paracasei* (06TCa22) isolated from the Mongolian dairy product. *International Immunopharmacology*, **17**(2): 358-365.
- Blaxhall, P.C. & Daisley, K.W. (1973). Routine haematological methods for use with fish blood. *Journal of Fish Biology*, **5**(6): 771-781.
- Bradford, M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, **72**(1-2): 248-254.

- Casillas, E. & Smith, L.S. (1977). Effect of stress on blood coagulation and haematology in rainbow trout (*Salmo gairdneri*). *Journal of Fish Biology*, **10**(5): 481-491.
- Cech Jr, J.J. & Wohlschlag, D.E. (1982). Seasonal patterns of respiration, gill ventilation, and hematological characteristics in the striped mullet, *Mugil cephalus* L. *Bulletin of Marine Science*, **32**(1): 130-138.
- Chakrabarty, P. & Banerjee, V. (1988). Effect of organophosphorus pesticides on the peripheral hemogram of the fish *Channa punctatus*. *Environment and Ecology*, **6**(2): 390-394.
- Chakrabarti, R. & Vasudeva, R.Y. (2006). *Achyranthes aspera* stimulates the immunity and enhances the antigen clearance in *Catla catla*. *International Immunopharmacology*, **6**(5): 782-790.
- Choi, S.H., Park, K.H., Yoon, T.J., Kim, J.B., Jang, Y.S. & Choe, C.H. (2008). Dietary Korean mistletoe enhances cellular non-specific immune responses and survival of Japanese eel (*Anguilla japonica*). *Fish & Shellfish Immunology*, **24**(1): 67-73.
- Citarasu, T. (2010). Herbal biomedicines: a new opportunity for aquaculture industry. *Aquaculture International*, **18**(3): 403-414.
- Cockson, A. & Bourne, D. (1972). Enzymes in the digestive tract of two species of euryhaline fish. *Comparative Biochemistry and Physiology--Part A: Physiology*, **41**(4): 715-718.
- Collazos, M.E., Ortega, E., Barriga, C. & Rodriguez, A.B. (1998). Seasonal variation in haematological parameters in male and female *Tinca tinca*. *Molecular and Cellular Biochemistry*, **183**(1-2): 165-168.
- Dalmo, R.A., Ingebrigtsen, K. & Bøgwald, J. (1997). Non-specific defence mechanisms in fish, with particular reference to the reticuloendothelial system (RES). *Journal of Fish Diseases*, **20**(4), 241-273.
- Erlanger, B.F., Kokowsky, N. & Cohen, W. (1961). The preparation and properties of two new chromogenic substrates of trypsin. *Archives of Biochemistry and Biophysics*, **95**(2): 271-278.
- Evelyn, T.P.T. (2002). Finfish immunology and its use in preventing infectious diseases in cultured finfish. *Diseases in Asian Aquaculture IV*, (CR Lavilla-Pitogo & ER Cruz-Lacierda ed.), 303-324.
- Farahi, A., Kasiri, M., Sudagar, M., Soleimani Iraei, M. & Zorriehzakra, S.M.J. (2012). Effect of dietary supplementation of *Melissa officinalis* and *Aloe vera* on hematological traits, lipid oxidation of carcass and performance in rainbow trout (*Oncorhynchus mykiss*). *Online Journal of Animal and Feed Research*, **1**: 1-5.
- Fukumoto, L. R. & Mazza, G. (2000). Assessing antioxidant and prooxidant activities of phenolic compounds. *Journal of Agricultural and Food Chemistry*, **48**(8): 3597-3604.
- Furne, M., Hidalgo, M.C., Lopez, A., Garcia-Gallego, M., Morales, A.E., Domezain, A., Domezaine, J. & Sanz, A. (2005). Digestive enzyme activities in Adriatic sturgeon *Acipenser naccarii* and rainbow trout *Oncorhynchus mykiss*. A comparative study. *Aquaculture*, **250**(1-2): 391-398.
- Garcia, M.P., Echevarria, G., Martinez, F.J. & Zamora, S. (1992). Influence of blood sample collection on the haematocrit value of two teleosts: rainbow trout (*Oncorhynchus mykiss*) and European sea bass (*Dicentrarchus labrax* L.). *Comparative biochemistry and physiology. Comparative Physiology*, **101**(4): 733-736.
- Gawlicka, A., Parent, B., Horn, M.H., Ross, N., Opstad, I. & Torrissen, O.J. (2000). Activity of digestive enzymes in yolk-sac larvae of Atlantic halibut (*Hippoglossus hippoglossus*): indication of readiness for first feeding. *Aquaculture*, **184**(3-4): 303-314.
- Gjedrem, T., Robinson, N. & Rye, M. (2012). The importance of selective breeding in aquaculture to meet future demands for animal protein: a review. *Aquaculture*, **350**: 117-129.
- Gudding, R., Lillehaug, A. & Evensen, Ø. (1999). Recent developments in fish vaccinology. *Veterinary Immunology and Immunopathology*, **72**(1-2): 203-212.
- Haghighi, M. & Rohani, M.S. (2013). The effects of powdered ginger (*Zingiber officinale*) on the haematological and immunological parameters of rainbow trout *Oncorhynchus mykiss*. *Journal of medicinal Plant and Herbal Therapy Research*, **1**(1): 8-12.
- Haghighi, M., Sharif Rohani, M., Samadi, M., Tavoli, M., Eslami, M. & Yusefi, R. (2014). Study of effects *Aloe vera* extract supplemented feed on hematological and immunological indices of rainbow trout (*Oncorhynchus mykiss*). *International Journal of Advanced Biological and Biomedical Research*, **2**(6): 2143-2154.
- Harikrishnan, R., Balasundaram, C. & Heo, M.S. (2011). Impact of plant products on innate and adaptive immune system of cultured finfish and shellfish. *Aquaculture*, **317**(1-4): 1-15.
- Harikrishnan, R., Jin, C.N., Kim, M.C., Kim, J.S., Balasundaram, C. & Heo, M.S. (2010). Effectiveness and immunomodulation of chemotherapeutants against scuticociliate *Philasterides dicentrarchi* in olive flounder. *Experimental Parasitology*, **124**(3): 306-314.
- Hickey, C.R. (1976). Fish hematology, its uses and significance. *NY Fish Game J*, **23**(2): 170-175.
- Ieri, F., Innocenti, M., Possieri, L., Gallori, S. & Mulinacci, N. (2015). Phenolic composition of "bud extracts" of *Ribes nigrum* L., *Rosa canina* L. and *Tilia tomentosa* M. *Journal of Pharmaceutical and Biomedical Analysis*, **115**: 1-9.
- Iqbal, K.J., Ashraf, M., Javid, A., Khan, N., Abbas, F., Hafeez-ur-Rehman, M., Rafique, M.K., Rasool, F., Azmat, H. & Altaf, M. (2016). Effect of Different Plant and Animal Origin (Fishmeal) Feeds on Digestive Enzyme Activity and Haematology of Juvenile *Labeo rohita*. *Pakistan Journal of Zoology*, **48**(1): 201-207.
- Iwama, G.K., Greer, G.L. & Larkin, P.A. (1976). Changes in some hematological characteristics of coho salmon (*Oncorhynchus kisutch*) in response to acute exposure to dehydroabietic acid (DHAA) at different exercise levels. *Journal of the Fisheries Board of Canada*, **33**(2): 285-289.
- Jian, J. & Wu, Z. (2004). Influences of traditional Chinese medicine on non-specific immunity of Jian carp (*Cyprinus carpio* var. Jian). *Fish & Shellfish Immunology*, **16**(2): 185-191.

- Johnston, J.R. (1978). Oxygen metabolism and the microbicidal activity of macrophages. *Federation Proceedings*, **37**(13): 2759-2764.
- Kaleeswaran, B., Ilavenil, S. & Ravikumar, S. (2011). Dietary supplementation with *Cynodon dactylon* (L.) enhances innate immunity and disease resistance of Indian major carp, *Catla catla* (Ham.). *Fish & Shellfish Immunology*, **31**(6): 953-962.
- Kawai, S. & Ikeda, S. (1973). Studies in digestive enzymes of fishes. III. Development of the digestive enzymes of rainbow trout after hatching and the effect of dietary change on the activities of digestive enzymes in the juvenile stage. *Bulletin of the Japanese Society for the Science of Fish*, **38**: 265-270.
- Khondoker, S., Hossain, M.M.M., Hasan-Uj-Jaman, M., Alam, M.E., Zaman, M.F.U. & Tabassum, N. (2016). Effect of *Nigella sativa* (Black Cumin Seed) to Enhance the Immunity of Common Carp (*Cyprinus carpio*) Against *Pseudomonas fluorescens*. *American Journal of Life Sciences*, **4**(3): 87-92.
- Kono, T., Hamasuna, S., Korenaga, H., Iizasa, T., Nagamine, R., Ida, T. & Sakai, M. (2012). The role of neuromedin U during inflammatory response in the common carp. *Fish & Shellfish Immunology*, **32**(1): 151-160.
- Kori-Siakpere, O., Ake, J.E.G. & Idoge, E. (2005). Haematological characteristics of the African snakehead, *Parachanna obscura*. *African Journal of Biotechnology*, **4**(6): 527-530.
- Kour, D., Sharma, L.L. & Sharma, B.K. (2004). Use of herb, Bala (*Sida cordifolia* Linn.) as growth promoter in the supplementary feed of *Cirrhinus mrigala* (Ham.). *Indian Journal of Fisheries*, **51**(4): 501-504.
- Kumar, I.V., Chelladurai, G., Veni, T., Peeran, S.S.H. & Mohanraj, J. (2014). Medicinal plants as immunostimulants for health management in Indian cat fish. *Journal of Coastal Life Medicine*, **2**(6): 426-430.
- Lee-Shing, F. & Shu-Fen, C. (1989). Effect of salinity on the activities of digestive proteases from the tilapia fish, *Oreochromis niloticus* in different culture environments. *Comparative Biochemistry and Physiology Part A: Physiology*, **93**(2): 439-443.
- Lewis, S.M., Bain, B.J., and Bates, I. (2001). Dacie and Lewis Practical Haematology, 9th ed. London: Churchill Livingstone; p 633.
- Mahdavi, M., Hajimoradloo, A. & Ghorbani, R., (2013). Effect of *Aloe vera* extract on growth parameters of common carp (*Cyprinus carpio*). *World Journal of Medical Sciences*, **9**: 55-60.
- Moriarty, D.J.W. (1973). The physiology of digestion of blue-green algae in the cichlid fish, *Tilapia nilotica*. *Journal of Zoology*, **171**(1): 25-39.
- Mostafa, A.A.Z.M., Ahmad, M.H., Mousallamy, A. & Samir, A. (2009). Effect of using dried Fenugreek seeds as natural feed additives on growth performance, feed utilization, whole-body coMyeloperoxidase activity sition and entropathogenic *Aeromonas hydrophila*-challenge of monsex Nile tilapia *O. niloticus* (L) fingerlings. *Australian Journal of Basic and Applied Sciences*, **3**(2): 1234-1245.
- Ojha, M.L., Chadha, N.K., Saini, V.P., Damroy, S. & Chandraprakash, S.P. (2014). Effect of ethanolic extract of *Mucuna pruriens* on growth, metabolism and immunity of *Labeo rohita* (Hamilton, 1822) fingerlings. *International Journal of Fauna & Biological Studies*, **1**(5): 1-9.
- Oluyemi, K.G., Adeprusi, E.A. & Olanrewage, J. (2008). Basic hematological parameters in African cat fish, *Clarias gariepinus* (Burchell, 1822) FED ascribe acid supplemented diets. *Research Journal of Animal Sciences*, **2**(1): 17-21.
- Örün, I., Dörücü, M. & Yazlak, H. (2003). Haematological parameters of three cyprinid fish species from Karakaya Dam Lake, Turkey. *OnLine Journal of Biological Sciences*, **3**: 320-328.
- Pakravan, S., Hajimoradloo, A. & Ghorbani, R. (2012). Effect of dietary willow herb, *Epilobium hirsutum* extract on growth performance, body composition, haematological parameters and *Aeromonas hydrophila* challenge on common carp, *Cyprinus carpio*. *Aquaculture Research*, **43**(6): 861-869.
- Patra, B.B., Panda, R.P. & Patra, A.K. (2014). Seasonal Variations in Certain Hematological Factors of *Catla catla* (Hamilton 1822). *Journal of Applied Physics*, **6**: 1-7.
- Rao, Y.V., Das, B.K., Jyotirmayee, P. & Chakrabarti, R. (2006). Effect of *Achyranthes aspera* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, **20**(3):, 263-273.
- Rodríguez, A., Esteban, M. & Meseguer, J. (2003). Phagocytosis and peroxidase release by seabream (*Sparus aurata* L.) leucocytes in response to yeast cells. *The Anatomical Record*, **272**(1): 415-423.
- Sahoo, P.K., Kumari, J. & Mishra, B.K. (2005). Non-specific immune responses in juveniles of Indian major carps. *Journal of Applied Ichthyology*, **21**(2): 151-155.
- Saurabh, S. & Sahoo, P.K. (2008). Lysozyme: an important defence molecule of fish innate immune system. *Aquaculture Research*, **39**(3): 223-239.
- Secombes, C.J. (1996). The nonspecific immune system: cellular defenses. *The Fish Immune System: Organism, Pathogen and Environment*, **15**: 63-103.
- Secombes, C.J. & Fletcher, T.C. (1992). The role of phagocytes in the protective mechanisms of fish. *Annual Review of Fish Diseases*, **2**: 53-71.
- Seung-Cheol, J.I., Jeong, G., Gwang-Soon, I.M., Lee, S., Yoo, J. & Takii, K. (2007). Dietary medicinal herbs improve growth performance, fatty acid utilization, and stress recovery of Japanese flounder. *Fisheries Science*, **73**(1): 70-76.
- Siddiqui, A.Q. & Naseem, S.M. (1979). The haematology of Rohu, *Labeo rohita*. *Journal of Fish Biology*, **14**(1): 67-72.
- Tekinay, A.A. & Davies, S.J. (2001). Dietary carbohydrate level influencing feed intake, nutrient utilisation and plasma glucose concentration in the rainbow trout, *Oncorhynchus mykiss*. *Journal of Veterinary and Animal Sciences*, **25**(5): 657-666.
- Worthington, C.S (1991). Worthington enzyme manual related Biochemical. Freehold, New Jersey, USA.

- Xie, J., Liu, B., Zhou, Q., Su, Y., He, Y., Pan, L. Ge, X. & Xu, P., (2008). Effects of anthraquinone extract from rhubarb *Rheum officinale* Bail on the crowding stress response and growth of common carp *Cyprinus carpio* var. *Jian*. *Aquaculture*, **281**: 5-11.
- Yin, G., Ardó, L., Thompson, K.D., Adams, A., Jeney, Z. & Jeney, G. (2009). Chinese herbs (*Astragalus radix* and *Ganoderma lucidum*) enhance immune response of carp, *Cyprinus carpio*, and protection against *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, **26**(1): 140-145.
- Yılmaz, S., Ergün, S. & Çelik, E.Ş. (2012). Effects of herbal supplements on growth performance of sea bass (*Dicentrarchus labrax*): Change in body composition and some blood parameters. *Journal of BioScience and Biotechnology*, **1**(3): 217-222.